





Loss of the Bacterial Flagellar Motor Switch Complex upon Cell Lysis

Mohammed Kaplan,^a  Elitza I. Tocheva,^{a*} Ariane Briegel,^{a*} Megan J. Dobro,^{a*} Yi-Wei Chang,^{a*} Poorna Subramanian,^a Alasdair W. McDowall,^a Morgan Beeby,^{a*}  Grant J. Jensen^{a,b}

^aDivision of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, USA

^bDepartment of Chemistry and Biochemistry, Brigham Young University, Provo, Utah, USA

ABSTRACT The bacterial flagellar motor is a complex macromolecular machine whose function and self-assembly present a fascinating puzzle for structural biologists. Here, we report that in diverse bacterial species, cell lysis leads to loss of the cytoplasmic switch complex and associated ATPase before other components of the motor. This loss may be prevented by the formation of a cytoplasmic vesicle around the complex. These observations suggest a relatively loose association of the switch complex with the rest of the flagellar machinery.

IMPORTANCE We show in eight different bacterial species (belonging to different phyla) that the flagellar motor loses its cytoplasmic switch complex upon cell lysis, while the rest of the flagellum remains attached to the cell body. This suggests an evolutionary conserved weak interaction between the switch complex and the rest of the flagellum which is important to understand how the motor evolved. In addition, this information is crucial for mimicking such nanomachines in the laboratory.

KEYWORDS cell lysis, flagellar motor, cryo-ET, switch complex

The bacterial flagellar motor is a complex nanomachine responsible for cell motility in a wide range of species. It consists of a cell envelope-embedded motor that rotates an extracellular filament, connected by a universal joint known as the hook (1). The motor is composed of a stator consisting of ion channels embedded in the cytoplasmic or inner membrane (IM) and a rotor consisting of a cytoplasmic switch complex (also called the C-ring), a membrane/supramembrane (MS)-ring, a periplasmic driveshaft known as the rod, associated bushings in the peptidoglycan cell wall (the P-ring), and, in diderms, the outer membrane (the lipopolysaccharide, or L-ring) (1). A flagellar type III secretion system (FT3SS) is responsible for self-assembly of the machine, which begins with the IM-associated components and proceeds in a stepwise fashion outward to the extracellular ones (1). In addition to this conserved core, the flagella of various species can have periplasmic or extracellular species-specific components that adorn their flagella (2, 3).

The flagellar machinery can also disassemble, as in the programmed ejection of the flagellum during the life cycle of the alphaproteobacterium *Caulobacter crescentus*. This process is thought to be accompanied by the digestion of the C terminus of the MS-ring protein FlIF (4). More recently, flagellar disassembly has been observed in many species under starvation or mechanical stress. This process starts with loss of flagellar hooks and filaments and continues with disassembly of motor components, leaving plugged P- and L-rings in the cell wall and outer membrane (5–11). Additionally, we have recently shown that programmed flagellar ejection in *C. crescentus* leaves similar plugged P- and L-rings suggesting an evolutionary link between this process and the starvation-induced one (12).

The complexity of the flagellar motor and its location spanning the cell envelope mean that it is challenging to purify intact. The development of cryogenic electron tomography

Citation Kaplan M, Tocheva EI, Briegel A, Dobro MJ, Chang Y-W, Subramanian P, McDowall AW, Beeby M, Jensen GJ. 2021. Loss of the bacterial flagellar motor switch complex upon cell lysis. *mBio* 12:e00298-21. <https://doi.org/10.1128/mBio.00298-21>.

Editor Gisela Storz, National Institute of Child Health and Human Development (NICHD)

Copyright © 2021 Kaplan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Grant J. Jensen, jensen@caltech.edu.

* Present address: Elitza I. Tocheva, Department of Microbiology and Immunology, Life Sciences Institute, The University of British Columbia, Vancouver, Canada; Ariane Briegel, Leiden University, Sylvius Laboratories, Leiden, The Netherlands; Megan J. Dobro, Hampshire College, Amherst, Massachusetts, USA; Yi-Wei Chang, Department of Biochemistry and Biophysics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA; Morgan Beeby, Department of Life Sciences, Imperial College London, South Kensington Campus, London, United Kingdom.

Received 5 February 2021

Accepted 26 April 2021

Published 8 June 2021




FIG 1 Slices through electron cryotomograms (except panels C, F, I, and L) of intact or lysed cells of various species showing flagella either with the switch complex (yellow ellipses) or lacking the switch complex (yellow arrows point to the expected position of the absent C-ring). Cyan ellipses indicate densities belonging to FlhA_c. Red boxes in panels N, Q, T, and W indicate regions enlarged in the panel to the right. (C, F, I, and L) Subtomogram averages (STA) of the subcomplexes shown in panels B, E, H, and K, respectively. Scale bars, 20 nm. IM, inner membrane; OM, outer membrane.

(cryo-ET), the highest-resolution imaging technique currently applicable to unique biological objects, has enabled the structure to be studied *in situ* in a native state inside cells. Over the past 15 years, our laboratory has collected tens of thousands of electron cryotomograms of dozens of bacterial species (13). This resource includes many examples of lysed cells, collected either accidentally or intentionally. For example, we have found that lysis of *Escherichia coli* cells by light penicillin treatment flattens them, making them more suitable for cryo-ET experiments (14). Similarly, the thick cell wall of *Bacillus subtilis* renders them a challenging sample for cryo-ET experiments, so we digested the cell wall of *B. subtilis* with lysozyme to visualize membrane-embedded structures in protoplasts and cell lysates. While examining these lysates, we identified a flagellar subcomplex that lacks the C-ring and contains only the MS-ring, the rod, and the hook (occasionally with part of the filament) (Fig. 1A and B). We identified a total of 71 such particles in 26 cryotomograms, enabling us to calculate a subtomogram average of the structure (Fig. 1C).

Wondering whether C-ring loss after cell lysis is a general phenomenon, we examined electron cryotomograms of lysed cells from different phyla (*Proteobacteria* and *Firmicutes*) available in our database. Besides the results in *B. subtilis*, we identified lysed cells retaining flagellar filaments in *E. coli* (penicillin treated), *Helicobacter hepaticus*, *Campylobacter jejuni*, *Hydrogenovibrio crunogenus*, *Hylemonella gracilis*, *Selenomonas* sp. (clinical isolate), and *Acetonebma longum*. In each case, in addition to fully intact flagella (which have the switch complex), we also observed flagellar filaments and hooks connected to motors lacking the C-ring (Fig. 1D to X). Multiple examples were found in some species that allowed us to average them (*H. hepaticus*, 7 examples in 7 lysed cells; *A. longum*, 7 examples in 2 lysed cells; *Selenomonas* sp., 6 examples in 2 lysed cells)




FIG 2 (A to G) Slices through electron cryotomograms of lysed cells of the indicated species, highlighting the presence of a flagellum with an intact C-ring (red arrows) encapsulated by an IM vesicle. Scale bars, 25 nm.

(Fig. 1F, I, and L). In roughly half of these partially disassembled motors, we observed densities corresponding to the C terminus of the FT3SS protein FlhA (FlhA_c) at the base of the MS-ring (Fig. 1H, K, O, R, and X, cyan circles).

Less frequently, lysed cells contained a small vesicle of IM encapsulating the cytoplasmic components of the flagellar motor (*C. jejuni*, 2 examples in 2 lysed cells; *A. longum*, 4 examples in 4 lysed cells; *H. hepaticus*, 4 examples in 4 lysed cells; single examples for the rest). These vesicles were only slightly larger than the switch complex, and, in each case, the C-ring was retained (Fig. 2).

As the cases mentioned above were either from complete cell lysates (as in *B. subtilis*), partially lysed cells (*E. coli*), or cells that were randomly lysed during sample preparation (the rest of the examples), we investigated this correlation between cell lysis and the loss of the switch complex by imaging cells that have undergone a more controlled lysis. To that end, we prepared and imaged *Treponema primitia* cells, which are spirochetes with periplasmic flagella, treated with different digestive enzymes: (i) cells incubated with 5 mg/ml lysozyme for 15 min prior to plunge-freezing, (ii) cells incubated with 5 mg/ml proteinase K for 15 min prior to plunge freezing, and (iii) undigested cells as a control group. The cells in groups 1 and 2 lacked an intact outer membrane and cell wall (Fig. 3). While the 25 motors identified in untreated cells all had C-rings, 6 of the 16 motors present in cells treated with lysozyme lacked the C-ring, and 1 motor out of the 9 motors in cells treated with proteinase K lacked the C-ring (Table 1, Fig. 3), suggesting again a correlation between cell lysis and C-ring loss.




FIG 3 Slices through electron cryotomograms of undigested (A and B), lysozyme-digested (D to G), or proteinase K-digested (H to K) *T. primitia* cells. Red boxes in panels A, D, F, H, and J indicate regions enlarged in the panel to the right. Flagella with or without the switch complex are indicated by yellow ellipses or yellow arrows (pointing to the expected position of the absent C-ring), respectively. Scale bars in panels A, D, F, H, and J, 100 nm; in panels B, E, G, I, and K, 50 nm. (C and L) Subtomogram averages (STA) of an intact flagellar motor (EMD-1235; see reference 31) (C) or a motor lacking the switch complex present in digested cells (L). Scale bars, 20 nm.

These results suggest that, upon cell lysis, the C-ring of some motors dissociates from the MS-ring but is retained if the proteins are encapsulated by a vesicle of cytoplasm. It is possible that the higher protease activity in the cytoplasm compared to the periplasm (15) plays a role in the loss of the cytoplasmic parts of the motor (i.e., the C-ring and the ATPase), but the preservation of the cytoplasmic FlhA_C suggests dissociation of the C-ring rather than proteolysis. This leads us to conclude that the interaction between the C- and MS-rings is more labile than interactions between other flagellar components. This also would explain the absence of the switch complex in early preparations of purified flagella (16, 17). Later studies found that the switch complex can be retained if flagella are purified under mild conditions of pH and salt (18–21), but

TABLE 1 Number of motors with or without C-ring in digested or untreated *T. primitia*

<i>Treponema primitia</i>	No. of motors with C-ring	No. of motors without C-ring	No. unclear	No. of tomograms
Untreated cells	25	0	0	18
+Lysozyme	8	6	2	13
+Proteinase K	8	1	0	5

even under mild conditions, the majority of purified flagella lose their C-rings after 20 minutes at room temperature or 80 hours at 4°C (20).

In our case, the lysis conditions were different for the various samples. For example, *B. subtilis* was completely lysed with lysozyme prior to plunge-freezing and imaging, and only cell lysate was present in the cryotomograms, while *E. coli* was mildly lysed by incubating the sample with penicillin before plunge-freezing. On the other hand, *T. primitia* was mildly lysed with lysozyme and proteinase K. The majority of the observed particles lacked the C-ring in the lysate of *B. subtilis*, but only a small proportion of motors were devoid of the switch complex in mildly lysed *E. coli* and *T. primitia*. While this could be related to the different lysis conditions and how severely the cells are lysed, we cannot exclude species-specific differences in this process. As for other species, the examples we identified were randomly lysed during sample preparation and we anticipate that the lysis conditions were mild (standard growth media), but we do not know how long individual cells were lysed before plunge-freezing. Whereas these results suggest that the loss of the switch complex is more prominent under severe lysis (like in *B. subtilis*), further experiments are required to quantify this process in relation to different lysis conditions (including mechanical stress) in different species.

In the case of partially lysed *T. primitia*, only the outer membrane and cell wall appeared damaged, while the cells continued to have an intact inner membrane and normal cytoplasm, as seen in our cryotomograms. It is known that the stators interact with the cell wall upon incorporation into the flagellar motor to get activated (22), and the damage of the cell wall in these cells could interfere with the incorporation of stators to the motor and their interaction with the switch complex. The loss of C-ring in these cells could be due to the interruption of this interaction, which would normally help to stabilize the C-ring in its place.

Previous studies of flagellar loss in intact *Vibrio alginolyticus* cells suggest that this process starts with the detachment, not digestion, of the C-ring, which is freed to diffuse along the inner membrane (9), and intact C-rings do not copurify with the motor in this species (23). Cryo-ET of intact *Shewanella oneidensis* cells lacking the filament proteins FlaA and FlaB ($\Delta flaAB$ mutant) also revealed a flagellar subcomplex with the extracellular hook and periplasmic components but without the cytoplasmic ones (6), further suggesting that C-ring loss is not limited to cell lysis. It is conceivable that releasing the C-ring is beneficial to cells in order to save energy when there is no filament to rotate, as in the *S. oneidensis* $\Delta flaAB$ mutant.

The phenomenon of breaking the bacterial flagellum and leaving stable flagellar subcomplexes under certain conditions is a recently described one. First, we and others showed that various bacteria can lose their flagella under starvation, leaving the P- and L-rings as a stable subcomplex in the outer membrane (5–11). However, how the motor breaks to leave PL-rings remains unknown. Using fluorescence microscopy, Zhuang et al. presented a model of the possibility of breaking the bacterial flagellum at the rod level, leaving the MS- and C-rings detached (9). While we have no evidence that our observations in the current work are connected to the starvation-related process, they add to the repertoire of stable flagellar subcomplexes under stress conditions, where the C-ring dissociates from the motor without the MS-ring leaving the rest of the flagellum (including the extracellular parts) as a stable structure.

The evolution of the bacterial flagellum is believed to have started as a primordial secretion system that subsequently added the periplasmic and extracellular components (rod, hook, and filament) by multiple gene diversification and duplication events (24). This connection between the flagellum and a secretion system is further bolstered by its structural similarity to the bacterial type III secretion system (25, 26), where it is believed that the evolution of the flagellum preceded that of the type III secretion system (27). However, a recent high-resolution structure of purified flagellar basal bodies highlighted structural differences between these two molecular machines (28). Presumably, tinkering of a primordial secretion system with ion channels and the addition of the switch complex led to the formation of the first motor. However, was the C-ring added before or

after the union between the preliminary secretion system and the ion channels? When did the rod, hook, and filament evolve? While both the preliminary secretion system and ion channels could have had independent functions unrelated to cell motility, it remains unknown what the role of the ancestral C-ring proteins was before functioning as a switch complex. Just as the secretion and assembly of the rod, hook, and filament proteins mirror their evolutionary past (24), it could be that the flagellar relics lacking the switch complex identified here, and the PL-subcomplexes in the starvation-related process, also reflect major modular transitions in the history of flagellar evolution.

Experimental procedures. Cell growth, sample preparation, cryo-ET imaging, and image processing were performed as previously described for *E. coli* and *H. hepaticus* (14), *H. crunogenus* (29), *A. longum* (30), *T. primitia* (31), and *Vibrio cholerae* (32). *Pseudomonas fluorescens* was grown in K10 medium as described in reference 33 and *C. jejuni* as described in reference 34. *H. gracilis* was grown for 48 h in broth 233 at 26°C without antibiotics to a final optical density at 600 nm of <0.1 and subsequently incubated with attack-phase *Bdellovibrio bacteriovorus* for 3 days, after which cells were spun down at 1,000 × *g* for 5 min and concentrated ~10× for plunge-freezing. *Selenomonas* spp. were isolated from the human gut by Emma Allen-Vercoe, University of Guelph. Cells were grown anaerobically in fastidious anaerobe agar supplemented with 5% defibrinated sheep's blood. *B. subtilis* protoplasts (obtained from the D. Kearns laboratory, Indiana University) were prepared with lysozyme using a modified protocol based on reference 35. Subsequent imaging and processing were performed as described for other species. Subtomogram averaging was done as in reference 6.

ACKNOWLEDGMENTS

This project was funded by the NIH (grant RO1 AI127401 to G.J.J.) and a Baxter postdoctoral fellowship from Caltech to M.K. Cryo-ET work was done in the Beckman Institute Resource Center for Transmission Electron Microscopy at the California Institute of Technology.

We are grateful to Catherine Oikonomou for critically reading the manuscript. We thank Cristina Iancu for collecting *H. crunogenus* data.

REFERENCES

- Macnab RM. 2003. How bacteria assemble flagella. *Annu Rev Microbiol* 57:77–100. <https://doi.org/10.1146/annurev.micro.57.030502.090832>.
- Chen S, Beeby M, Murphy GE, Leadbetter JR, Hendrixson DR, Briegel A, Li Z, Shi J, Tocheva EI, Müller A, Dobro MJ, Jensen GJ. 2011. Structural diversity of bacterial flagellar motors: structural diversity of bacterial flagellar motors. *EMBO J* 30:2972–2981. <https://doi.org/10.1038/emboj.2011.186>.
- Kaplan M, Ghosal D, Subramanian P, Oikonomou CM, Kjaer A, Pirbadian S, Ortega DR, Briegel A, El-Naggar MY, Jensen GJ. 2019. The presence and absence of periplasmic rings in bacterial flagellar motors correlates with stator type. *Elife* 8:e43487. <https://doi.org/10.7554/eLife.43487>.
- Jenal U, Shapiro L. 1996. Cell cycle-controlled proteolysis of a flagellar motor protein that is asymmetrically distributed in the *Caulobacter* predivisional cell. *EMBO J* 15:2393–2406. <https://doi.org/10.1002/j.1460-2075.1996.tb00597.x>.
- Ferreira JL, Gao FZ, Rossmann FM, Nans A, Brenzinger S, Hosseini R, Wilson A, Briegel A, Thormann KM, Rosenthal PB, Beeby M. 2019. γ -Proteobacteria eject their polar flagella under nutrient depletion, retaining flagellar motor relic structures. *PLoS Biol* 17:e3000165. <https://doi.org/10.1371/journal.pbio.3000165>.
- Kaplan M, Subramanian P, Ghosal D, Oikonomou CM, Pirbadian S, Starwalt-Lee R, Magewaran SK, Ortega DR, Gralnick JA, El-Naggar MY, Jensen GJ. 2019. *In situ* imaging of the bacterial flagellar motor disassembly and assembly processes. *EMBO J* 38:e100957. <https://doi.org/10.15252/embj.2018100957>.
- Zhu S, Schniederbernd M, Zhitnitsky D, Jain R, Galán JE, Kazmierczak BI, Liu J. 2019. *In situ* structures of polar and lateral flagella revealed by cryo-electron tomography. *J Bacteriol* 201:e00117–19. <https://doi.org/10.1128/JB.00117-19>.
- Kaplan M, Sweredoski MJ, Rodrigues J, Tocheva EI, Chang Y-W, Ortega DR, Beeby M, Jensen GJ. 2020. Bacterial flagellar motor PL-ring disassembly subcomplexes are widespread and ancient. *Proc Natl Acad Sci U S A* 117:8941–8947. <https://doi.org/10.1073/pnas.1916935117>.
- Zhuang X, Guo S, Li Z, Zhao Z, Kojima S, Homma M, Wang P, Lo C, Bai F. 2020. Live-cell fluorescence imaging reveals dynamic production and loss of bacterial flagella. *Mol Microbiol* 114:279–291. <https://doi.org/10.1111/mmi.14511>.
- Zhu S, Gao B. 2020. Bacterial flagella loss under starvation. *Trends Microbiol* 28:785–788. <https://doi.org/10.1016/j.tim.2020.05.002>.
- Zhuang X-Y, Lo C-J. 2020. Construction and loss of bacterial flagellar filaments. *Biomolecules* 10:1528. <https://doi.org/10.3390/biom10111528>.
- Kaplan M, Wang Y, Chreifi G, Zhang L, Chang Y-W, Jensen GJ. 2021. Programmed flagellar ejection in *Caulobacter crescentus* leaves PL-subcomplexes. *J Mol Biol* 433(13):167004. <https://doi.org/10.1016/j.jmb.2021.167004>.
- Ding HJ, Oikonomou CM, Jensen GJ. 2015. The Caltech tomography database and automatic processing pipeline. *J Struct Biol* 192:279–286. <https://doi.org/10.1016/j.jsb.2015.06.016>.
- Briegel A, Ortega DR, Tocheva EI, Wuichet K, Li Z, Chen S, Muller A, Iancu CV, Murphy GE, Dobro MJ, Zhulin IB, Jensen GJ. 2009. Universal architecture of bacterial chemoreceptor arrays. *Proc Natl Acad Sci U S A* 106:17181–17186. <https://doi.org/10.1073/pnas.0905181106>.
- Talmadge K, Gilbert W. 1982. Cellular location affects protein stability in *Escherichia coli*. *Proc Natl Acad Sci U S A* 79:1830–1833. <https://doi.org/10.1073/pnas.79.6.1830>.
- DePamphilis ML, Adler J. 1971. Purification of intact flagella from *Escherichia coli* and *Bacillus subtilis*. *J Bacteriol* 105:376–383. <https://doi.org/10.1128/JB.105.1.376-383.1971>.
- Aizawa SI, Dean GE, Jones CJ, Macnab RM, Yamaguchi S. 1985. Purification and characterization of the flagellar hook-basal body complex of *Salmonella typhimurium*. *J Bacteriol* 161:836–849. <https://doi.org/10.1128/JB.161.3.836-849.1985>.

18. Driks A, DeRosier DJ. 1990. Additional structures associated with bacterial flagellar basal body. *J Mol Biol* 211:669–672. [https://doi.org/10.1016/0022-2836\(90\)90063-R](https://doi.org/10.1016/0022-2836(90)90063-R).
19. Schuster SC, Baeuerlein E. 1992. Location of the basal disk and a ringlike cytoplasmic structure, two additional structures of the flagellar apparatus of *Wolinella succinogenes*. *J Bacteriol* 174:263–268. <https://doi.org/10.1128/JB.174.1.263-268.1992>.
20. Khan IH, Reese TS, Khan S. 1992. The cytoplasmic component of the bacterial flagellar motor. *Proc Natl Acad Sci U S A* 89:5956–5960. <https://doi.org/10.1073/pnas.89.13.5956>.
21. Francis NR, Sosinsky GE, Thomas D, DeRosier DJ. 1994. Isolation, characterization and structure of bacterial flagellar motors containing the switch complex. *J Mol Biol* 235:1261–1270. <https://doi.org/10.1006/jmbi.1994.1079>.
22. Hosking ER, Vogt C, Bakker EP, Manson MD. 2006. The *Escherichia coli* MotAB proton channel unplugged. *J Mol Biol* 364:921–937. <https://doi.org/10.1016/j.jmb.2006.09.035>.
23. Koike M, Terashima H, Kojima S, Homma M. 2010. Isolation of basal bodies with C-ring components from the Na⁺-driven flagellar motor of *Vibrio alginolyticus*. *J Bacteriol* 192:375–378. <https://doi.org/10.1128/JB.01121-09>.
24. Liu R, Ochman H. 2007. Stepwise formation of the bacterial flagellar system. *Proc Natl Acad Sci U S A* 104:7116–7121. <https://doi.org/10.1073/pnas.0700266104>.
25. Blocker A, Komoriya K, Aizawa S-I. 2003. Type III secretion systems and bacterial flagella: insights into their function from structural similarities. *Proc Natl Acad Sci U S A* 100:3027–3030. <https://doi.org/10.1073/pnas.0535335100>.
26. Diepold A, Armitage JP. 2015. Type III secretion systems: the bacterial flagellum and the injectisome. *Philos Trans R Soc B Biol Sci* 370:20150020. <https://doi.org/10.1098/rstb.2015.0020>.
27. Abby SS, Rocha EPC. 2012. The non-flagellar type III secretion system evolved from the bacterial flagellum and diversified into host-cell adapted systems. *PLoS Genet* 8(9):e1002983. <https://doi.org/10.1371/journal.pgen.1002983>.
28. Tan J, Zhang X, Wang X, Xu C, Chang S, Wu H, Wang T, Liang H, Gao H, Zhou Y, Zhu Y. 2021. Structural basis of assembly and torque transmission of the bacterial flagellar motor. *Cell* 184(10):2665–2679.e19. <https://doi.org/10.1016/j.cell.2021.03.057>.
29. Iancu CV, Morris DM, Dou Z, Heinhorst S, Cannon GC, Jensen GJ. 2010. Organization, structure, and assembly of α -carboxysomes determined by electron cryotomography of intact cells. *J Mol Biol* 396:105–117. <https://doi.org/10.1016/j.jmb.2009.11.019>.
30. Tocheva EI, Dekas AE, McGlynn SE, Morris D, Orphan VJ, Jensen GJ. 2013. Polyphosphate storage during sporulation in the gram-negative bacterium *Acetonebacterium longum*. *J Bacteriol* 195:3940–3946. <https://doi.org/10.1128/JB.00712-13>.
31. Murphy GE, Leadbetter JR, Jensen GJ. 2006. In situ structure of the complete *Treponema primitia* flagellar motor. *Nature* 442:1062–1064. <https://doi.org/10.1038/nature05015>.
32. Yang W, Alvarado A, Glatter T, Ringgaard S, Briegel A. 2018. Baseplate variability of *Vibrio cholerae* chemoreceptor arrays. *Proc Natl Acad Sci U S A* 115:13365–13370. <https://doi.org/10.1073/pnas.1811931115>.
33. Boyd CD, Smith TJ, El-Kirat-Chatel S, Newell PD, Dufrene YF, O'Toole GA. 2014. Structural features of the *Pseudomonas fluorescens* biofilm adhesin LapA required for LapG-dependent cleavage, biofilm formation, and cell surface localization. *J Bacteriol* 196:2775–2788. <https://doi.org/10.1128/JB.01629-14>.
34. Müller A, Beeby M, McDowall AW, Chow J, Jensen GJ, Clemons WM. 2014. Ultrastructure and complex polar architecture of the human pathogen *Campylobacter jejuni*. *MicrobiologyOpen* 3:702–710. <https://doi.org/10.1002/mbo3.200>.
35. DeCastro-Costa MR, Landman OE. 1977. Inhibitory protein controls the reversion of protoplasts and L forms of *Bacillus subtilis* to the walled state. *J Bacteriol* 129:678–689. <https://doi.org/10.1128/JB.129.2.678-689.1977>.